

## Cervical cancer screening, women 18+

Health Priority A: Access to Primary and Preventive Health Services

Objective A2b: Increase provider screening for chronic diseases and other health risks.

2010 Target: No target established

Indicator: Cervical cancer screening, women 18+

### Percent of Wisconsin Women Aged 18+ Who Had a Pap Smear in the Past Three Years

Year (N)	Total	
	Percent	+/-
2000	87%	2%
(N)	1,197	
2001	90%	1%
(N)	1,535	
2002	88%	1%
(N)	1,925	
2004	86%	2%
(N)	2,058	

Source: Wisconsin Behavioral Risk Factor Survey, Bureau of Health Information and Policy, Division of Public Health, Department of Health and Family Services.

Note: Excludes women who reported having had a hysterectomy.

N        Number in sample. See data documentation.

+/-      Confidence interval. See data documentation.

**Continued on next page**

Prepared for Tracking the State Health Plan, 2010—State-Level Data

<http://dhfs.wisconsin.gov/statehealthplan/track2010> (July 2005)

Bureau of Health Information and Policy

Division of Public Health

Wisconsin Department of Health and Family Services

## Cervical cancer screening, women 18+, continued

### Percent of Wisconsin Women Aged 18+ Who Had a Pap Smear in the Past Three Years, by Race/Ethnicity

Years (N)	Total		African American*		Hispanic		White*	
	Percent	+/-	Percent	+/-	Percent	+/-	Percent	+/-
2000-2002	88%	1%	92%	3%	87%	5%	88%	1%
(N)	4,657		432		180		3,860	
2001-2002, 2004	88%	1%	89%	6%	83%	8%	88%	1%
(N)	5,530		554		207		4,567	

Source: Wisconsin Behavioral Risk Factor Survey, Bureau of Health Information and Policy, Division of Public Health, Department of Health and Family Services.

Note: Excludes women who reported having had a hysterectomy.

N        Number in sample. See data documentation.

+/-      Confidence interval. See data documentation.

\*Non-Hispanic